

FAST FACT AND CONCEPT #41
PRESSURE ULCERS: DEBRIDEMENT AND DRESSINGS

Charles von Gunten, MD and Frank Ferris, MD

Background *Fast Fact #40* discussed the staging and prevention of pressure ulcers; this *Fast Fact* discusses their management. The first step in deciding how to manage pressure ulcers is an assessment of whether the wound is likely to heal. If the patient has a prognosis of months to years, adequate nutrition, and blood flow to the tissue, then healing is possible. If the patient has a prognosis of days to weeks, anorexia/cachexia, and/or the wound has inadequate perfusion, then symptom control alone is appropriate and uncomfortable/burdensome treatments are not appropriate.

Debridement Always provide adequate analgesia! Necrotic tissue must be removed for ulcer healing; surgical debridement is the fastest and most effective method when there is healthy surrounding tissue. Note: If the patient is close to dying, and/or the wound will never heal, then debridement should not be attempted. Debridement gels (such as Hypergel, Santyl, Nu-gel) are applied onto an ulcer under an occlusive dressing (such as DuoDerm), are available for ulcers that do not require surgery or when surgical debridement is incomplete. These products come with or without enzymes to encourage autolytic or enzymatic debridement. For minimally necrotic ulcers, occlusive dressings such as DuoDerm, changed weekly, promote autolysis.

A commonly prescribed form of mechanical debridement is the use of saline, wet-dry dressings. This treatment retards healing by pulling off new epithelial cells as part of healthy granulation tissue; its use for the treatment of skin ulcers should be abandoned.

Antimicrobials Ulcer healing is delayed if there is bacterial infection within the wound bed. Erythema, purulent exudate, and fever are signs of infection. Cleansing and application of topical antibiotics may be sufficient for superficial infection with minimal surrounding erythema. Systemic antibiotics are indicated for deep/surrounding tissue infection, or if ulcer healing is delayed. Cleanse wounds that are expected to heal with non-cytotoxic fluids (e.g., saline). Cytotoxic fluids (e.g., Betadine) will kill granulation tissue. **Clinical Pearl:** do not cleanse an ulcer with any fluid you would not put in your eye if you want the ulcer to heal.

Dressings Living tissue requires moisture for transport of oxygen and nutrients. A moist ulcer environment promotes the migration of fibroblasts and epithelial cells; growth factors are present in the serous exudate that speed healing. In contrast, a dry environment is conducive to necrosis and eschar.

There are 6 classes of dressings distinguished by the wear time and whether you want to add or remove fluid to maintain the ideal moist, interactive ulcer-healing environment. A dry ulcer needs to have moisture added through a hypotonic gel (donates water). With wet exudates, a hypertonic gel or foam is used to remove water.

1. *Polyurethane foams* (LYOfoam, Allevyn, Nu-Derm, Flexzan): most absorptive; used under a covering secondary dressing.
2. *Alginates* (Kaltostat, Sorbsan): desiccate an overly wet wound, prevent maceration of surrounding skin from excess fluid, and are hemostatic and may reduce infection risk.
3. *Hydrogels* (IntraSite, Elasto-Gel, ClearSite, Aquasorb): used for dry, abrasions, radiation skin damage, minor burns. Likely should be changed at least every four days. May require a secondary dressing to secure.
4. *Hydrocolloid wafers* (DuoDerm, Comfeel, Tegasorb, Restore): promote autolysis, angiogenesis, and granulation. Self-adhesive. Remain in place for 5-7 days. Often used to "seal" a wound that is otherwise clean to promote healing. Can also be used to seal an underlying dressing to maintain a moist environment in which the wound can heal. Note: do not to use an occlusive dressing if there is a substantial risk of infection.
5. *Thin films* (OpSite, Tegaderm): for skin at risk or Stage I pressure ulcers. Can also hold another type of absorbent dressing in place.

6. *Cotton Gauze*: used to cover the primary dressing. Rarely an appropriate dressing for a significant skin ulcer. Note: Saline wet-to-dry dressings are only useful for mechanical debridement.

References

1. Walker P. The pathophysiology and management of pressure ulcers. In: *Topics in Palliative Care, Volume 3*. Portenoy RK, Bruera E, eds. New York, NY: Oxford University Press; 1998: pp 253-270.
2. Walker P. Update on pressure ulcers. *Principles & Practice of Supportive Oncology Updates*. 2000; 3(6):1-11.

Version History: This *Fast Fact* was originally edited by David E Weissman MD. 2nd Edition published August 2005; 3rd edition May 2015. Current version re-copy-edited March 2009; May 2015; and again in April 2021.

Fast Facts and Concepts are edited by Sean Marks MD (Medical College of Wisconsin) and associate editor Drew A Rosielle MD (University of Minnesota Medical School), with the generous support of a volunteer peer-review editorial board, and are made available online by the [Palliative Care Network of Wisconsin](#) (PCNOW) and the Center to Advance Palliative Care (www.capc.org). *Fast Facts and Concepts* are editorially independent of PCNOW and the Center to Advance Palliative Care, and the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact's* content. The full set of *Fast Facts* are available at <http://www.mypcnow.org/#!/fast-facts/cb1h> or <http://www.capc.org/fast-facts/> along with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-NonCommercial 4.0 International Copyright (<http://creativecommons.org/licenses/by-nc/4.0/>). *Fast Facts* can only be copied and distributed for non-commercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment and consult other relevant and up-to-date experts and resources. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.