

FAST FACTS AND CONCEPTS #84 SWALLOW STUDIES, TUBE FEEDING, AND THE DEATH SPIRAL

David E Weissman MD

Introduction The reflex by families and doctors to provide nutrition for the patient who cannot swallow is overwhelming. It is now common practice for such patients to undergo a swallowing evaluation and if there is significant impairment to move forward with feeding tube placement (either nasogastric or gastrostomy) – see *Fast Fact #128*. Data suggest that in-hospital mortality for hospitalizations in which a feeding tube is placed is 15-25%, and one year mortality after feeding tube placement is 60%. Predictors of early mortality include: advanced age, CNS pathology (stroke, dementia), cancer (except early stage head/neck cancer), disorientation, and low serum albumin.

The Tube Feeding Death Spiral The clinical scenario, the *tube feeding death spiral*, typically goes like this:

1. Hospital admission for complication of “brain failure” or other predictable end organ failure due to primary illnesses (e.g. urosepsis in setting of advanced dementia).
2. Inability to swallow and/or direct evidence of aspiration and/or weight loss with little oral intake.
3. Swallowing evaluation followed by a recommendation for non-oral feeding either due to aspiration or inadequate intake.
4. Feeding tube placed leading to increasing “agitation” leading to patient-removal or dislodgement of feeding tube.
5. Re-insertion of feeding tube; hand and/or chest restraints placed.
6. Aspiration pneumonia.
7. Intravenous antibiotics and pulse oximetry.
8. Repeat 4 – 6 one or more times.
9. Family conference.
10. Death.

Note: at my institution, the finding of a dying patient with a feeding tube, restraints, and pulse oximetry is known as *Weissman’s triad*.

Suggestions

- Recognize that the inability to maintain nutrition through the oral route, in the setting of a chronic life-limiting illness and declining function, is usually *a marker of the dying process*. Discuss this with families as a means to a larger discussion of overall end of life goals.
- Ensure that your colleagues are aware of the key data and recommendations on tube feedings (see *Fast Fact #10*).
- Ensure there is true informed consent prior to feeding tube insertion—families *must* be given alternatives (e.g. hand feeding, comfort measures) along with discussion of goals and prognosis.
- Assist families by providing information and a clear recommendation for or against the use of a feeding tube. Families who decide against feeding tube placement can be expected to second guess their decision and will need continued team support.
- If a feeding tube is placed establish clear goals (e.g. improved function) and establish a timeline for re-evaluation to determine if goals are being met (typically 2-4 weeks).

References

1. Finucane TE, et al. Tube feeding in patients with advanced dementia. *JAMA*. 1999; 282:1365-1369.

2. Finucane TE, Bynum JP. Use of tube feeding to prevent aspiration pneumonia. *Lancet*. 1996; 348:1421-24.
3. Cowen ME et al. Survival estimates for patients with abnormal swallowing studies. *JGIM*. 1997; 12:88-94.
4. Rabeneck L, et al. Long term outcomes of patients receiving percutaneous endoscopic gastrostomy tubes. *JGIM*. 1996; 11:287-293.
5. Grant MD, et al. Gastrostomy placement and mortality among hospitalized Medicare beneficiaries. *JAMA*. 1998; 279:1973-1976.
6. Mitchell SL. Clinical Crossroads: a 93-year-old man with advanced dementia and eating problems. *JAMA*. 2007; 298:2527-2536.
7. Cervo FA, Bryan L, Farber S. To PEG or not to PEG. A review of evidence for placing feeding tubes in advanced dementia and the decision-making process. *Geriatrics*. 2006; 61:30-35.

Version History: This *Fast Fact* was originally edited by David E Weissman MD. 2nd Edition was edited by Drew A Rosielle MD and published October 2007; 3rd Edition June 2015.

Fast Facts and Concepts are edited by Sean Marks MD (Medical College of Wisconsin) and associate editor Drew A Rosielle MD (University of Minnesota Medical School), with the generous support of a volunteer peer-review editorial board, and are made available online by the [Palliative Care Network of Wisconsin](#) (PCNOW); the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact*'s content. The full set of *Fast Facts* are available at [Palliative Care Network of Wisconsin](#) with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-NonCommercial 4.0 International Copyright (<http://creativecommons.org/licenses/by-nc/4.0/>). *Fast Facts* can only be copied and distributed for non-commercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment and consult other relevant and up-to-date experts and resources. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.