

wFAST FACTS AND CONCEPTS #104
INSOMNIA: NON-PHARMACOLOGIC TREATMENTS
Robert Arnold MD, Michael Miller, and Rohtesh Mehta, MD MPH

Introduction This *Fast Fact* addresses non-pharmacologic therapies for insomnia; *Fast Fact* #101 discusses patient assessment and #105 discusses pharmacologic treatment of insomnia.

Non-Specific Treatments

I. **Improving Sleep Hygiene** Sleep hygiene education should be provided to anyone with insomnia. First, there is no arbitrary rule for how much someone should sleep. Generally, older patients need less sleep. Second, patients should be advised to keep a regular sleep schedule; this means going to bed and getting up at the same time. Third, patients should avoid long daytime naps, alcohol, and caffeine. These principles may be difficult for seriously ill patients, particularly maintaining a regular routine for hospitalized patients and avoiding naps for fatigued, seriously ill patients.

II. **Behavioral Treatments**

- Relaxation therapies: The patient can be taught to use various relaxation techniques just prior to bedtime such as progressive muscle relaxation, guided imagery, meditation, diaphragmatic breathing or hypnosis (see *Fast Fact* #211).
- Stimulus control therapy: This focuses on establishing a connection between the bed and sleep. It emphasizes not watching TV, reading in bed, or using bedroom for any other activity (except sexual activity).
- Sleep restriction therapy: This therapy requires patient motivation as it involves restricting the time allowed in bed depending upon the actual time spent in bed over the past 2 weeks and then going to bed 15 minutes earlier every night (increased weekly) until the target bedtime is achieved. Sleep compression therapy is a variant of sleep restriction but the reduction in sleep is made gradually.
- Cognitive behavioral therapy: This is a multi-component psychotherapy, and includes sleep restriction, stimulus control and cognitive therapy which focuses on identifying unwanted feelings or thoughts and replacing them with more positive thoughts. It is effective in treating chronic insomnia in the general population, in patients with chronic pain, and in women with metastatic breast cancer.

The last two therapies are the only evidence-based psychological treatments for insomnia in older adults.

III. **Physical Exercises**: Although the data is limited, studies in elderly patients suggest significantly improved sleep duration and onset latency and minimal enhancement in sleep efficiency with community-based endurance training programs such as low impact aerobics, brisk walking, and Tai Chi Chih.

Choosing which therapy to use first depends mainly on provider experience and patient motivation. It is difficult to know how long non-pharmacologic therapy should be tried before attempting other measures. This depends on the patient, the severity of the insomnia, or the severity of the illness. Many studies which showed cognitive behavioral therapy to be effective used a treatment regimen of 7-8 weeks.

Specific Treatments *Obstructive sleep apnea* is treated with positive airway pressure (e.g., 'CPAP') ventilation at night. Although some patients report difficulty becoming accustomed to sleeping with the CPAP mask on, this therapy can dramatically improve symptoms. Surgery is sometimes indicated for obstructive sleep apnea. *Symptoms* from an underlying medical disorder can also contribute to insomnia. Adequately treating a patient's pain, nausea and vomiting (See *Fast Fact* #5), or dyspnea (See *Fast Fact* #27) should improve sleep.

Spiritual and existential concerns can be an important cause of insomnia in palliative care patients. Patients may be able to avoid these concerns during the day through the distraction of

daily activities but have difficulty ignoring them at night. Thus, it is important to directly address a patient's spiritual concerns, worries, and fears about dying during the day. Brief psychotherapy may be helpful.

References

1. Montgomery P, Dennis JA. Cognitive behavioural interventions for sleep problems in adults aged 60+. *Cochrane Database of Systematic Reviews* 2003, Issue 1. Art. No.: CD003161. DOI: 10.1002/14651858.CD003161.
2. Montgomery P, Dennis JA. Physical exercise for sleep problems in adults aged 60+. *Cochrane Database of Systematic Reviews* 2002, Issue 4. Art. No.: CD003404. DOI: 10.1002/14651858.CD003404.
3. McCurry SM, Logsdon RG, Teri L, Vitiello MV. Evidence-based psychological treatments for insomnia in older adults. *Psychol Aging*. 2007;22(1):18-27
4. Quesnel C, et al. Efficacy of cognitive-behavioral therapy for insomnia in women treated for nonmetastatic breast cancer. *J Consulting Clinical Psych*. 2003; 71(1):189-200.
5. Jungquist CR, O'Brien C, Matteson-Rusby S, et al. The efficacy of cognitive-behavioral therapy for insomnia in patients with chronic pain. *Sleep Med*. 2010 Feb 2
6. Irwin MR, Olmstead R, Motivala SJ. Improving sleep quality in older adults with moderate sleep complaints: A randomized controlled trial of Tai Chi Chih. *Sleep*. 2008;31(7):1001-8.

Author Affiliations: University of Pittsburgh Medical Center, Pittsburgh, PA.

Version History: This *Fast Fact* was originally written by Michael Miller and Robert Arnold, edited by David E Weissman MD, and published in November 2003. Re-copy-edited in April 2009. 2nd Edition was updated by Robert Arnold and Rohtesh Mehta and published in September 2011. Revised in July 2013 to correct an error (that CPAP is the primary positive airway pressure treatment for OSA, not bilevel ventilation).

Fast Facts and Concepts are edited by Sean Marks MD (Medical College of Wisconsin) and associate editor Drew A Rosielle MD (University of Minnesota Medical School), with the generous support of a volunteer peer-review editorial board, and are made available online by the [Palliative Care Network of Wisconsin](#) (PCNOW); the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact's* content. The full set of *Fast Facts* are available at [Palliative Care Network of Wisconsin](#) with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-NonCommercial 4.0 International Copyright (<http://creativecommons.org/licenses/by-nc/4.0/>). *Fast Facts* can only be copied and distributed for non-commercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment and consult other relevant and up-to-date experts and resources. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.