

**FAST FACTS AND CONCEPTS #121
ORAL MUCOSITIS: DIAGNOSIS AND ASSESSMENT**

Clarissa F Henson and Robert Arnold MD

Background The term mucositis refers to the inflammatory response of the oral-pharyngeal mucosa resulting from systemic chemotherapy or from radiotherapy that includes the oral-pharyngeal mucosa within the radiation field. The term stomatotoxicity is specific to mucositis affecting the oral mucosa. Mucositis results from the destruction of rapidly dividing epithelial cells of the oral-pharyngeal mucosal epithelium and the secondary release of inflammatory mediators such as TNF-alpha and interleukin-1 beta. This *Fast Fact* discusses the diagnosis and assessment of mucositis; *Fast Fact* #130 discusses its prevention and treatment.

The Radiation Therapy Oncology Group (RTOG) describes five grades of acute mucositis:

- Grade 1: Injection; may experience mild pain not requiring analgesics.
- Grade 2: Patchy mucositis which may produce an inflammatory serosanguinitis discharge; may experience moderate pain requiring analgesia.
- Grade 3: Confluent fibrinous mucositis; may include severe pain requiring opioid analgesics.
- Grade 4: Ulceration, hemorrhage or necrosis.
- Grade 5: Death resulting from mucositis.

Causes Both patient-related factors and treatment-related factors influence the severity of mucositis. Increased total dose of radiation, fraction size, and volume of normal tissue in the irradiated field all increase the risk of mucositis. Not all chemotherapy agents produce the same risk of mucositis; 5-fluoruracil, doxorubicin and methotrexate commonly cause mucositis while vincristine does not. The simultaneous combination of radiation and chemotherapy used in head and neck cancer will cause more intense mucositis than single-modality therapy. Patient-related factors such as the overall condition of the oral mucosa prior to therapy, pre-existing xerostomia, pre-existing collagen-vascular disorders, the underlying nutritional status, and the development of neutropenia during therapy all impact the development and severity of mucositis.

In head and neck cancer, virtually all patients undergoing radiation, with or without chemotherapy, will develop grade 1 and 2 mucositis. More severe mucositis (grade 3 or higher) develops in approximately 41% of patients receiving combined radiation and chemotherapy to the head and neck and in 21% of patients receiving radiation therapy alone.

Clinical Findings & Natural History Clinical signs of mucosal damage and cell death appear after the first 1 to 2 weeks of radiation therapy and as early as 3 days after chemotherapy. Initial mucosal damage results in patchy erythema, edema, atrophy and whitening of the mucosal tissue with increased sensitivity—patients report a burning sensation in the mouth. Further loss of mucosal epithelium becomes most prominent in the fourth or fifth week of standard fractionation radiation resulting in fibrinous exudation, confluent inflammation, and ulceration (see *Fast Facts* # 66, 67). The mouth sores and swelling can lead to significant oral pain, pain with swallowing, weight loss and dehydration. Mucosal damage predisposes the patient to oral superinfection leading to further pain and alterations in taste and appetite and decreased quality of life. Resolution of oral mucositis occurs several weeks after the completion of RT and chemotherapy.

Assessment A thorough assessment of the patient with mucositis should include:

- Pain assessment to include thorough review of analgesics used, effect and toxicities.
- Nutritional assessment.
- Quality of life assessment including screening for depression.
- Complete oral examination; assessment for local fungal (see *Fast Fact* #147), bacterial, or viral infections.

References

1. Bernier J, Dommé C, Ozsahin M, et al. Postoperative irradiation with or without concomitant chemotherapy for locally advanced head and neck cancer. *N Engl J Med*. 2004; 350:1945-1952.
2. Cooper JS, Pajak TF, Forastiere AA, et al. Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *N Engl J Med*. 2004; 350:1937-1944.
3. Fu K, et al. A Radiation Therapy Oncology Group (RTOG) phase III randomized study to compare hyperfractionation and two variants of accelerated fractionation to standard fractionation radiotherapy for head and neck squamous cell carcinomas: first report of RTOG 9003. *Int J Radiat Oncol Biol Phys*. 2000; 48:7-16.

Version History: This *Fast Fact* was originally edited by David E Weissman MD and published in September 2004. Version re-copy-edited in April 2009; revised again July 2015.

Fast Facts and Concepts are edited by Sean Marks MD (Medical College of Wisconsin) and associate editor Drew A Rosielle MD (University of Minnesota Medical School), with the generous support of a volunteer peer-review editorial board, and are made available online by the [Palliative Care Network of Wisconsin](#) (PCNOW); the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact's* content. The full set of *Fast Facts* are available at [Palliative Care Network of Wisconsin](#) with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-NonCommercial 4.0 International Copyright (<http://creativecommons.org/licenses/by-nc/4.0/>). *Fast Facts* can only be copied and distributed for non-commercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment and consult other relevant and up-to-date experts and resources. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.