

FAST FACTS AND CONCEPTS #429
DE-STIGMATIZING THE LANGUAGE OF ADDICTION

Janet Ho MD MPH, Katie F Jones MSN APN, Zachary Sager MD, Sarah Wakeman MD, Jessica S. Merlin MD PhD

Background Discrimination and stigmatization of patients who use psychoactive drugs or who have addiction is common in healthcare (1–4). Using people-first, medically accurate, and stereotype-reducing language with colleagues, trainees, and in documentation (5), is fundamental to dispelling stigma (6) and misperceptions, such as the myth of addiction as a willful, personal choice or moral failing. The right words can convey empathy and reduce mistrust in healthcare for patients who have often faced a history of marginalization. This *Fast Facts* discusses healthcare stigmatization of patients who use drugs or have addiction, and will present preferred language that respects personhood, supports recovery, and promotes accurate knowledge about the full spectrum of substance use and an understanding of addiction as a chronic, treatable condition (see *Fast Fact* #127).

Stigma in healthcare Stigma is a social process that occurs in the context of power (7), characterized by labeling, stereotyping, and discrimination based on real or perceived attributes. Stigma towards people who use drugs or develop addiction, as well as medications to treat this condition, are long-standing. Opioid use disorder (OUD) has long been treated apart from mainstream medicine in the US, from the 1914 Harrison Narcotics Act which separated treatment of OUD away from mainstream healthcare (8), to the racist “War on Drugs” which treated drug use as a criminal-legal issue removed from public health. Clinicians themselves commonly have negative biases about addiction (9–11) which is often reinforced during medical training via stigmatizing language (12). For example, patients described as “substance abusers” were perceived by clinicians as more threatening, more worthy of blame, and less likely to benefit from treatment compared with “persons with an SUD” (13). Altogether, stigma worsens the mental and physical health of people who use drugs (14–16), disproportionately affects communities of color (17), and harms patients on long-term therapeutic opioids (18,19). It even underlies the crisis of overdose deaths, as only 12.5% of people with OUD receive medication proven to reduce accidental overdose (18,19). The table below presents preferred language for clinicians.

Preferred language (20,21)	Non-preferred language
<i>Substance use disorder (SUD), opioid use disorder (OUD), alcohol use disorder (AUD), unhealthy/risky use (preferred to misuse), non-medical use, addiction (which can be used to mean severe SUD per the DSM 5)</i>	<i>Substance/drug/alcohol abuse, drug/drinking problem.</i> The terms <i>dependence</i> and <i>addiction</i> are both used by patients; however, <i>dependence</i> clinically refers to physiologic withdrawal reactions when a substance is stopped. It may occur without a SUD and should not be used synonymously by clinicians with SUD/addiction.
<i>Addiction is a chronic condition characterized by the compulsive use of a substance despite harmful consequences.</i>	Using language like <i>choice, lifestyle, moral failing, lack of will power, or personal failure</i> to describe SUD/addiction.
<i>Person with a SUD, person with addiction, person who uses drugs, person who injects drugs (PWID)</i>	<i>Addict, drug/substance abuser, person with a drug habit, alcoholic, IV drug user, drug-seeker</i>
<i>Person not actively using, person in remission from SUD, person in recovery</i>	<i>Clean, former addict.</i> The term <i>sober</i> is generally not used by clinicians, though some patients use it themselves.
<i>Substance present/not present in urine screen</i>	<i>Dirty/clean urine.</i> Describing urine as <i>positive</i> or <i>negative</i> for a substance can be clinically confusing.
<i>Medications for OUD (MOUD), Medication for addiction treatment (MAT), Opioid agonist therapy (OAT)</i>	<i>Opioid replacement therapy and opioid substitution</i> are not preferred as they stigmatize MOUD as ‘replacing one addiction with another.’ <i>Medication assisted treatment</i> is

	not preferred because MOUD alone can save lives without additional treatment for some patients.
<i>Patient-directed discharge, patient left before treatment completion despite medical recommendations, patient declined further treatment after informed consent</i>	<i>Leaving against medical advice, AMA, eloped:</i> these clinical designations have no clinical, regulatory, or professional standards. As such they can stigmatize patients, worsen health outcomes, and disproportionately affect vulnerable populations (22)
<i>Undertreated pain, risky opioid use (e.g., self-titration of meds, etc.) related to undertreated pain (instead of pseudoaddiction) Using opioids to treat non-pain symptoms (instead of chemical coping)</i>	<i>Pseudoaddiction, chemical coping:</i> these non-diagnostic euphemisms are applied inconsistently and subject to unconscious bias by, for example, clinicians to describe patients with OUD, for whom the clinician does not feel comfortable 'labeling' as having OUD. This stigmatizes OUD and can prevent appropriate treatment (23)
<i>Opioids</i>	<i>Narcotics</i> (which is used in a legal context to refer to multiple classes of illegal substances not just opioids)

Summary The language used to describe substance use and addiction must convey respect for individuals, an accurate understanding of the full spectrum of substance use (from non-problematic use to chaotic use and addiction), and effective treatment. Avoid perpetuating negative stereotypes by first recognizing, addressing, and combatting misunderstandings and stigma.

Authors' Affiliations: University of California San Francisco, San Francisco, CA (JH); VA Boston Healthcare System, Boston, MA (KFJ, ZS); New England Geriatric Research Education and Clinical Center, Boston, MA (ZS); Massachusetts General Hospital, Boston, MA (SW); University of Pittsburgh, Pittsburgh, PA (JM).

Conflicts of Interest: The authors have declared no conflicts of interest.

Version History: originally edited by Drew A Rosielle MD; first electronically published September 2021

References

1. Volkow ND. Stigma and the Toll of Addiction. *N Engl J Med.* 2020 Apr 2;382(14):1289–90.
2. Woods JS, Joseph H. Stigma from the Viewpoint of the Patient. *J Addict Dis.* 2015 Jul 3;34(2–3):238–47.
3. Ebenau A, Dijkstra B, ter Huurne C, Hasselaar J, Vissers K, Groot M. Palliative care for patients with substance use disorder and multiple problems: a qualitative study on experiences of healthcare professionals, volunteers and experts-by-experience. *BMC Palliat Care.* 2020 Dec;19(1):8.
4. McCradden MD, Vasileva D, Orchanian-Cheff A, Buchman DZ. Ambiguous identities of drugs and people: A scoping review of opioid-related stigma. *Intl J Drug Policy.* 2019 Dec 1;74:205–15.
5. P. Goddu A, O'Connor KJ, Lanzkron S, Saheed MO, Saha S, Peek ME, et al. Do Words Matter? Stigmatizing Language and the Transmission of Bias in the Medical Record. *J Gen Intern Med.* 2018 May;33(5):685–91.
6. Wakeman SE. Language and Addiction: Choosing Words Wisely. *Am J Public Health.* 2013 Apr;103(4):e1–2.
7. Link BG, Phelan JC. Conceptualizing Stigma. *Ann Rev Sociology.* 2001;27(1):363–85.
8. Courtwright DT. Preventing and Treating Narcotic Addiction--Century of Federal Drug Control. *N Engl J Med.* 2015 Nov 26;373(22):2095–7.
9. Adams VJM, Volkow ND. Ethical Imperatives to Overcome Stigma Against People With Substance Use Disorders. *AMA Journal of Ethics.* 2020 Aug 1;22(8):702–8.
10. van Boekel LC, Brouwers EPM, van Weeghel J, Garretsen HFL. Stigma among health professionals towards patients with substance use disorders and its consequences for healthcare delivery: Systematic review. *Drug Alcohol Dep.* 2013 Jul 1;131(1):23–35.
11. Atkins J, Dopp AL, Temaner and EB. Combatting the Stigma of Addiction— The Need for a Comprehensive Health System Approach. *NAM Perspectives [Internet].* 2020 Nov 9. Available from: <https://nam.edu/combating-the-stigma-of-addiction-the-need-for-a-comprehensive-health-system-approach/>. Accessed August 23, 2021.

12. Avery J, Knoepfmacher D, Mauer E, et al. Improvement in Residents' Attitudes Toward Individuals with Substance Use Disorders Following an Online Training Module on Stigma. *HSS Jnl*. 2019 Feb 1;15(1):31–6.
13. Kelly JF, Dow SJ, Westerhoff C. Does our choice of substance-related terms influence perceptions of treatment need? An empirical investigation with two commonly used terms. *J Drug Issues*. 2010;40(4):805–18.
14. Bearnot B, Mitton JA, Hayden M, Park ER. Experiences of care among individuals with opioid use disorder-associated endocarditis and their healthcare providers: Results from a qualitative study. *J Subst Abuse Treatment*. 2019 Jul 1;102:16–22.
15. Simon R, Snow R, Wakeman S. Understanding why patients with substance use disorders leave the hospital against medical advice: A qualitative study. *Subst Abus*. 2020;41(4):519–25.
16. Wakeman SE, Rich JD. Barriers to Medications for Addiction Treatment: How Stigma Kills. *Substance Use & Misuse*. 2018 Jan 28;53(2):330–3.
17. Race and the Drug War. Drug Policy Alliance [Internet]. Available from: <https://drugpolicy.org/issues/race-and-drug-war>. Accessed Aug, 23 2021.
18. Wakeman SE, Larochelle MR, Ameli O, et al. Comparative Effectiveness of Different Treatment Pathways for Opioid Use Disorder. *JAMA Netw Open*. 2020 Feb 5;3(2):e1920622–e1920622.
19. Tsai AC, Kiang MV, Barnett ML, et al. Stigma as a fundamental hindrance to the United States opioid overdose crisis response. *PLOS Medicine*. 2019 Nov 26;16(11):e1002969.
20. Kelly JF, Wakeman SE, Saitz R. Stop Talking 'Dirty': Clinicians, Language, and Quality of Care for the Leading Cause of Preventable Death in the United States. *Am J Med*. 2015 Jan 1;128(1):8–9.
21. Botticelli MP, Koh HK. Changing the Language of Addiction. *JAMA*. 2016 Oct 4;316(13):1361–2.
22. Alfandre D, Brenner J, Onukwugha E. Things We Do For No Reason: Against Medical Advice Discharges. *J Hosp Med*. 2017 Oct 2;12(10):843–5.
23. Bell K, Salmon A. Pain, physical dependence and pseudoaddiction: Redefining addiction for 'nice' people? *Intl J Drug Poicy*. 2009 Mar 1;20(2):170–8.

Fast Facts and Concepts are edited by Sean Marks MD (Medical College of Wisconsin) and associate editor Drew A Rosielle MD (University of Minnesota Medical School), with the generous support of a volunteer peer-review editorial board, and are made available online by the [Palliative Care Network of Wisconsin](#) (PCNOW); the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact's* content. The full set of *Fast Facts* are available at [Palliative Care Network of Wisconsin](#) with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-NonCommercial 4.0 International Copyright (<http://creativecommons.org/licenses/by-nc/4.0/>). *Fast Facts* can only be copied and distributed for non-commercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment and consult other relevant and up-to-date experts and resources. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.